

# Hessian fly resistance genes *H16* and *H17* are mapped to a resistance gene cluster in the distal region of chromosome 1AS in wheat

Lingrang Kong · Sue E. Cambron ·  
Herbert W. Ohm

Received: 23 March 2007 / Accepted: 22 June 2007 / Published online: 11 July 2007  
© Springer Science+Business Media B.V. 2007

**Abstract** Hessian fly [*Mayetiola destructor* (Say)] is one of the major insect pests of wheat (*Triticum aestivum* L.) worldwide. Hessian fly (Hf)-resistance genes *H16* and *H17* were reported to condition resistance to Hf biotype L that is prevalent in many wheat-growing areas of eastern USA, and both of them were previously assigned to wheat chromosome 5A by their linkage to *H9*. The objectives in this study were to (1) map *H16* and *H17* independent of their linkage with *H9* and (2) identify DNA markers that co-segregate with *H16* or *H17*, and that are useful for selection of these genes in segregating populations and to combine these genes with other Hf-resistance genes in wheat cultivars. Contrary to previously reported locations, *H16* and *H17* did not show linkage with the molecular markers on chromosome 5A. Instead, both of them are linked with the molecular markers on the short arm of chromosome 1A (1AS). The simple sequence repeat (SSR) marker *XpSP2999* and EST-derived SSR (eSSR)

marker *Xwem6b* are two flanking markers that are linked to *H16* at genetic distances of 3.7 and 5.5 cM, respectively. Similarly, *H17* is located between markers *XpSP2999* and *Xwem6b* at genetic distances of 6.2 and 5.1 cM, respectively. Five other SSR and eSSR markers including *Xcfa2153*, *Xbarc263*, *Xwem3a*, *Xwmc329*, and *Xwmc24* were also linked to *H16* and *H17* at close genetic distances. These closely linked molecular markers should be useful for pyramiding *H16* and *H17* with other Hessian fly resistance genes in a single wheat genotype. In addition, using Chinese Spring deletion line bin mapping we positioned all of the linked markers and the Hf-resistance genes (*H16* and *H17*) to the distal 14% of chromosome 1AS, where Hf-resistance genes *H9*, *H10*, and *H11* are located. Our results together with previous studies suggest that Hf-resistance genes *H9*, *H10*, *H11*, *H16*, and *H17* along with the pathogen resistance genes *Pm3* and *Lr10* appear to occupy a resistance gene cluster in the distal region of chromosome 1AS in wheat.

Contribution from Purdue Univ. Agric. Res. Programs Journal Article No. 2007-18105.

L. Kong · H. W. Ohm (✉)  
Department of Agronomy, Purdue University,  
915 W. State St, West Lafayette, IN 47907, USA  
e-mail: hohm@purdue.edu

S. E. Cambron  
Crop Production and Pest Control Research Unit  
USDA-ARS-MWA and Department of Entomology,  
Purdue University, West Lafayette, IN 47907, USA

**Keywords** *Triticum aestivum* · Microsatellite · eSSR · Gene mapping · Hessian fly resistance · Marker-assisted selection (MAS)

## Introduction

Genes in wheat that confer resistance to the Hessian fly (Hf) provide the most efficient and economical means of crop protection against this damaging insect

(Berzonsky et al. 2003). Currently, at least 32 Hf-resistance genes have been identified in wheat and its wild relatives, and these resistance genes have been designated in a series from *H1* to *H32* (Delibes et al. 1997; Ratcliffe and Hatchett 1997; McIntosh et al. 2003; Williams et al. 2003; Martín-Sánchez et al. 2003; Sardesai et al. 2005; Liu et al. 2005a, b, c; Wang et al. 2006; Zhao et al. 2006). Many of these genes, including *H6*, *H9–H11*, *H14–H20*, *H28*, *H29* and *H31*, were identified in tetraploid durum wheat, *Triticum turgidum* (AABB,  $2n = 4x = 28$ ) ssp. *Durum* Desf., and some of these genes, including *H13*, *H22*, *H23*, *H24*, *H26*, and *H32*, originated from *Aegilops tauschii* (DD,  $2n = 2x = 14$ ) Coss. Gene *H6* was located on chromosome 5A by monosomic analysis (Gallun and Patterson 1977). Genes *H3* and *H9* were shown by segregation analysis to be linked to *H6*, and *H15* was shown to be closely linked or allelic to *H9*, composing the linkage block *H3–H6–H9–H15* (Patterson and Gallun 1977; Stebbins et al. 1982; Maas et al. 1989). In addition, *H16* and *H17* were both assigned to wheat chromosome 5A, showing linkage with *H9* and/or *H10* (Obanni et al. 1988; Patterson et al. 1988; Ohm et al. 1995). Genes *H16* and *H17* confer resistance against Hessian fly biotype L, the most virulent and prevalent biotype in eastern USA.

Molecular mapping technologies provide new tools to map genes precisely on chromosomes of wheat. Ma et al. (1994) identified restriction fragment length polymorphism (RFLP) markers that are linked to *H23* and *H24* on 6D and 3DL, respectively. *H25* was mapped on 6BL with RFLP markers (Delaney et al. 1995). Williams et al. (2003) mapped *H31* on chromosome 5BS by its linkage to a sequence tagged site (STS) marker and an amplified fragment length polymorphism (AFLP). Recently, simple sequence repeat (SSR) or microsatellite markers have greatly facilitated Hf-resistance gene mapping. *H9*, *H10* and *H11* were mapped near the distal end of chromosome 1AS (Kong et al. 2005; Liu et al. 2005a) and not on 5A, as had been previously reported (Stebbins et al. 1980, 1982). *H13* was recently mapped on the distal region of chromosome 6DS (Liu et al. 2005c) and not 6DL, as was previously reported (Gill et al. 1987). *H26* was recently mapped on chromosome 3D, rather than 4D (Wang et al. 2006). In addition, *H22* was mapped to the short arm of chromosome 1DS (Zhao et al.

2006) and *H32* was mapped to 3DL (Sardesai et al. 2005), respectively, using SSR and other PCR-based DNA markers.

The genetic interaction between *Triticum* and *Mayetiola destructor* has been established as a gene-for-gene relationship (Gallun and Hatchett 1968; Hatchett and Gallun 1970). During compatible interactions with developing seedlings, Hf larvae establish feeding sites among the leaf sheaths near the crown at the base of the plant. Larvae induce the nutritive tissue 2–3 days after initiating their attack (Harris et al. 2006) and feed for 10–11 days (Gagné and Hatchett 1989; Harris et al. 2003; Anderson and Harris 2006). Feeding by virulent larvae not only stunts plant growth but also causes seedlings to accumulate more chloroplasts (Cartwright et al. 1959; Robinson et al. 1960). The irreversible changes in wheat tissues create a satisfactory environment to allow larval growth by delivering nutrients and protecting the larvae among the stunted leaf sheaths. Wheat seedlings without an effective R gene die or produce few seeds (Anderson and Harris 2006). During incompatible interactions, Hf larval feeding triggers activation of wheat genes in defense (Williams et al. 2002; Giovanini et al. 2006, 2007). However, the first-instar larvae do not die immediately. Rather, they continue to probe and move down the plant for 3–4 days (Gallun 1977; Grover 1995) and are unable to establish permanent feeding sites (Grover 1995). Larvae do not appear to grow on resistant host genotypes and typically die within 2–5 days after attempting to feed near the base of the plant (Harris et al. 2003; Anderson and Harris 2006; Giovanini et al. 2007). This incompatible interaction appears to have little impact on the plant because infested, resistant wheat plants exhibit relatively normal growth (Gallun et al. 1961).

Continuous evolution of virulent Hf genotypes necessitates the deployment of new resistance genes to prevent crop losses due to Hf infestation. Sequential deployment of single resistance genes has been suggested as an efficient strategy for crop protection from Hf (Cox and Hatchett 1986; Smith et al. 1999). Gould (1986) predicted that the resistance of a cultivar containing multiple genes for resistance to a single biotype of the Hf could be effective up to 20 times longer than resistance of a cultivar with a single resistance gene. The development of DNA markers that co-segregate with specific resistance genes and

that are efficient to use for plant selection, greatly enhance the feasibility of combining two or more Hf-resistance genes for deployment in wheat cultivars (Williams et al. 2003).

The objectives of this study were to: (i) map genes *H16* and *H17*, independent of their association with *H9*, and (ii) identify markers that co-segregate with *H16* or *H17* and that are useful in future MAS in wheat breeding for Hf resistance.

## Materials and methods

### Hessian fly stocks

Hf biotype L (virulent to resistance genes *H3*, *H5*, *H6*, and *H7H8*) is maintained by the USDA-ARS Crop Production and Pest Control Research Unit at Purdue University, in a 4°C cold room as a purified laboratory stock. Hf pupae together with infested susceptible wheat plants were stored at 4°C until Hf adults were needed.

### Plant materials

The wheat (*T. aestivum* L.) parent lines used for this study consisted of cultivar Len, which is susceptible to all known biotypes of the Hf; and Hf-resistant parent lines P921682 (*H16H16*) and P921680 (*H17H17*). The corresponding Hf-resistant durum wheat (*T. turgidum*) source lines were PI 94587 for *H16* and PI 428435 for *H17*. Hf resistance gene, *H16*, was transferred to the susceptible durum wheat line D6647 by backcrossing, D6647\*2/PI 94587, and testing the BC<sub>1</sub> progeny to Hf biotype L (Patterson et al. 1988). A resulting *H16H16* line was crossed to susceptible cultivar Newton, after which six cycles of backcrossing to Newton were carried out with selection for Hf biotype L-resistant plants after each cycle of backcrossing, followed by three generations of self-pollination and testing to biotype L to identify wheat line P921682 (*H16H16*). Similarly, PI 428435 was first backcrossed to D6647, identifying gene *H17* (Obanni et al. 1988) and resulting in a *H17H17* durum line. Gene *H17* was subsequently transferred from the *H17H17* durum line into common wheat cultivar Newton by backcrossing to Newton and selection for Hf biotype L-resistant plants after each of six cycles of backcrossing followed by three

generations of self-pollination and testing to biotype L, resulting in the wheat line P921680 (*H17H17*).

*Triticum aestivum* wheat line Chinese Spring (CS) and CS deletion lines del1AS-1 (KSU#4510-1) and del1AS-3 (KSU#4510-3) were included for mapping in this study.

### Progeny screening for Hf resistance

The Hf-resistant parental lines, P921682 and P921680, were, respectively, crossed to the susceptible *T. aestivum* cultivar Len, and the resulting F<sub>1</sub> plants were backcrossed to Len to produce BC<sub>1</sub>F<sub>1</sub> plant populations of 113 and 103 plants, respectively, from Len\*2/P921682 and Len\*2/P921680. BC<sub>1</sub>F<sub>1</sub> plants were self-pollinated to produce BC<sub>1</sub>F<sub>2</sub> seeds. Additionally, F<sub>2</sub> populations were derived from the crosses Len/P921682 and Len/P921680, respectively; and F<sub>2</sub> plants were self-pollinated to produce F<sub>3</sub> seeds. Plant tissue for DNA extraction was collected from all BC<sub>1</sub>F<sub>1</sub> and F<sub>2</sub> plants. BC<sub>1</sub>F<sub>2</sub> and F<sub>2:3</sub> families, along with parent lines Len, P921682, P921680 and resistance source lines PI 94587 and PI 428435 were scored for resistance to Hf biotype L.

Parent lines, F<sub>1</sub> plants and F<sub>2:3</sub> families, and BC<sub>1</sub>F<sub>2</sub> families, together with the resistance source lines, were seeded in soil in wooden flats that were 54 × 36 × 8 cm deep, containing 10 evenly spaced rows (Ohm et al. 1995). All Hf resistance screens were conducted as progeny tests. At least 15 seedlings of each BC<sub>1</sub>F<sub>2</sub> and F<sub>2:3</sub> family were grown in nine of the 10 rows. One center row was divided to accommodate both resistant and susceptible parental lines of 10 seedlings each as checks. Ten seedlings of each F<sub>1</sub> and the two durum resistance source lines were also included in the tests.

Hf bioassays were conducted similar to that described previously (Ohm et al. 1995; Anderson and Harris 2006). Briefly, plants were infested with Hf adults at the one-leaf stage by allowing the Hf to emerge from infested wheat plants under a cheese-cloth tent which covered the flats. Mating and oviposition proceeded for 24 h when all adult flies were removed. After infestation, flats were moved to growth chambers maintained at constant temperature (18 ± 1°C) with a 12-h light period.

Three weeks after infestation, F<sub>2:3</sub> and BC<sub>1</sub>F<sub>2</sub> families as well as F<sub>1</sub>, parents and the resistance source lines were classified as resistant, susceptible or

segregating. Susceptible plants appeared stunted, with dark green leaves, and live larvae at the base of the first leaf. Resistant plants appeared normal. For families that were segregating, the non-stunted plants were dissected to distinguish non-infested escapes from resistant plants by verifying the presence of dead red larvae at the base of the leaf sheath as proof that the seedling was infested and resistant. Hf resistance was evaluated as described by Ohm et al. (1995).

#### DNA isolation

Genomic DNA was isolated from seedling leaves using the CTAB method described by Saghai-Marooof et al. (1984) with minor modifications. A 1.67% CTAB extraction buffer [100 mM Tris–HCl buffer pH 8.0, 1.67 % (w/v) hexadecyltrimethylammonium bromide (CTAB), 100 mM Na<sub>2</sub>EDTA, and 1.4 M NaCl] was used. DNA concentration was quantified on a Hoefer DyNA Quant 200 Fluorometer (Hoefer Pharmacia Biotech Inc., Dubuque, IA, USA).

#### Bulked segregant analysis

For bulked segregant analysis (BSA) (Michelmore et al. 1991), equivalent amounts of genomic DNA from seven resistant (all of their tested F<sub>2:3</sub> progeny were resistant) and seven susceptible F<sub>2</sub> plants (all of their tested F<sub>2:3</sub> progeny were susceptible) from each population derived from the crosses Len/P921682 and Len/P921680, were, respectively, pooled to form resistant and susceptible bulks as described in Kong et al. (2005). Both bulks were used along with the parents to identify markers showing polymorphisms between the four samples. These polymorphic markers were further used to analyze BC<sub>1</sub>F<sub>1</sub> populations to determine linkages between SSR/eSSR markers and resistance genes *H16* and *H17*, respectively.

#### Microsatellite and STS marker analysis

Wheat microsatellite markers on chromosome 1A and 5A designated as either *Xgwm* for Gatersleben (Germany) wheat microsatellite (Röder et al. 1998) or *Xgdm* for Gatersleben D-genome microsatellite (Pestsova et al. 2000), or *Xwmc* for Wheat Microsatellite Consortium (Gupta et al. 2002) were tested for useful polymorphisms. Additional markers including

*Xbarc* (Beltsville Agriculture Research Center), *Xksm* (Kansas State University microsatellite), *Xcni* (Cornell University microsatellite), *Xcfa* and *Xcfd* (Pierre Sourdille microsatellite), *Xpsp* (Devos et al. 1995; Stephenson et al. 1998) on chromosome 1A were also tested on the *H16* and *H17* populations from the crosses Len/P921682 and Len/P921680, respectively. Considering that both *Pm3* and *Lr10* are located on chromosome 1AS, STS marker for *Pm3* (*STS-Pm3*, forward primer 5' ATGGCTAGATGCCCGTTATG 3' and reverse primer 5' AGAGCAGAGCAGTGCAACAA 3') and the STS marker for *Lr10* (*STS-Lr10*, forward primer 5' GCGCTATGCCTAACCTGAAG 3' and reverse primer 5' CTCCACATAGGCAGCACTGA 3') were developed based on the available sequences from the GenBank database (GenBank #AY605285 and GenBank #AY270157, respectively).

#### Wheat EST-derived microsatellite marker analysis

The Perl script MISA (<http://www.pgrc.ipk-gatersleben.de/misa.html>) was used to identify SSRs in a wheat EST database containing approximately one-half million ESTs (<http://www.wheat.pw.usda.gov/cgi-bin/ace/search/wEST>). Twenty-one EST-derived microsatellites (eSSRs) were chosen from the short arms of chromosomes from group 1 (Table 1) using seven consensus maps (Peng and Lapitan 2005). All of the primers were designed by Primer3 (<http://www.frodo.wi.mit.edu/cgi-bin/primer3/primer3>) and synthesized by Integrated DNA Technologies (Coralville, IA, USA).

#### Polymerase chain reaction

Polymerase chain reaction (PCR) for each SSR and STS marker was performed in a Bio-Rad MyCycler<sup>TM</sup>. Thermal Cycler (Hercules, CA, USA) at amplifications of 94°C for 2 min, followed by 35 cycles of 94°C for 30 s 50°C, 52°C, 55°C, or 60°C (based on primer annealing temperature) for 40 s, and 72°C for 1 min, with a final extension at 72°C for 7 min before cooling to 4°C. Each PCR (25 µl) consisted of 40 ng of template DNA, 10 mM Tris HCl, pH 8.3, 50 mM KCl, 0.1% Triton X-100, 1.5 mM MgCl<sub>2</sub>, 200 µM of each dNTP, 0.25 µM of each primer and 1 unit of *Taq* DNA polymerase. The amplified PCR products were

**Table 1** eSSR primers targeting short arms of wheat group 1 chromosomes applied to linkage analysis of Hf-resistance genes *H16* and *H17*

Locus <sup>a</sup>	Left primer	Right primer	EST accession	SSR motif	Tm <sup>b</sup> (°C)	Amplicon size (bp)	Chromosome location
<i>Xwem3a</i>	GATCTGTGACCGAGGCAGA	GCTGTGGAGGTCCAAAATGT	BG607867	(AC)7	55	77	1AS
<i>Xwem6a</i>	CCTGCTCTGCCATTACTTGG	TGCACCTCCATCTCCTTCTT	BF483588	(AG)12	55	145	IDS
<i>Xwem6b</i>	CCTGCTCTGCCATTACTTGG	TGCACCTCCATCTCCTTCTT	BF483588	(AG)12	55	150	1AS
<i>Xwem6c</i>	CCTGCTCTGCCATTACTTGG	TGCACCTCCATCTCCTTCTT	BF483588	(AG)12	55	165	IBS
<i>Xwem7f</i>	ACGGCGTGTGAGTTTTTCT	CAACTGCAACAACAAAACAGT	BE500104	(T)10	55	328	IBS
<i>Xwem8b</i>	TGTGCTTCAAGCCTCAAGTG	GCTCGCACTCGAGTACACTG	BE443007	(TAC)5	55	132	IDS
<i>Xwem9a</i>	CACCATCACCGAGATCCAA	GGAGCTCTCTCCACCTTGTG	BE494877	(CAGG)5	55	78	IBS
<i>Xwem9b</i>	CACCATCACCGAGATCCAA	GGAGCTCTCTCCACCTTGTG	BE494877	(CAGG)5	55	115	IBS
<i>Xwem9g</i>	CACCATCACCGAGATCCAA	GGAGCTCTCTCCACCTTGTG	BE494877	(CAGG)5	55	190	1AS
<i>Xwem10</i>	GAACATTTTGGCTCTGTG	TGGTGATCCAGAAAGCCATT	BF483804	(A)11	50	68	IDS
<i>Xwem11c</i>	CAGAGCAACAGATGTTGGA	TGCACGTAGTAGTAGGCACCTC	BG314086	(ACT)5	60	300	IDS
<i>Xwem11d</i>	CAGAGCAACAGATGTTGGA	TGCACGTAGTAGTAGGCACCTC	BG314086	(ACT)5	60	555	IBS
<i>Xwem12a</i>	CAGCAACCAATTACCAACACA	GCGAAAAATGATGGTTGTTGA	BE424439	(ACA)5	60	100	IDS
<i>Xwem12b</i>	CAGCAACCAATTACCAACACA	GCGAAAAATGATGGTTGTTGA	BE424439	(ACA)5	60	140	1AS
<i>Xwem12c</i>	CAGCAACCAATTACCAACACA	GCGAAAAATGATGGTTGTTGA	BE424439	(ACA)5	60	153	IBS
<i>Xwem20d</i>	GACACCTTCTCTTGCTCCAAA	GAAAGACGTGATCAGCATGGA	BE442801	(TTG)5	50	230	1AS
<i>Xwem20f</i>	GACACCTTCTCTTGCTCCAAA	GAAAGACGTGATCAGCATGGA	BE442801	(TTG)5	50	580	1AS
<i>Xwem25a</i>	CGCCTCAGAGCTCTTCACC	AAGATACGGTCCGTGTAGGAG	BG607143	(CCG)5	55	110	IBS
<i>Xwem46a</i>	ACGTTGTCTCCGTGTCAATTG	GGTCATGGCCCTCAGTCTCA	BE500430	(TCC)5	50	105	IDS
<i>Xwem47c</i>	CCTTCTCGACTCCCTCTTTCG	CCATTGCTCTGGACCTGT	BE426787	(AGG)5	55	600	IBS
<i>Xwem54c</i>	AGCCAAAGGAGCTGGAGGAC	GGCTCCGTGCTCTCTCGAC	BE517937	(CCG)5	55	370	IBS

<sup>a</sup> Wem = Wheat EST-derived microsatellite<sup>b</sup> Tm = melting temperature (°C)

fractionated on 2.0–3.0% agarose gels (based on the size difference of the polymorphism) using a mixture of 1:1 Metaphor<sup>®</sup> and Seakem<sup>®</sup> in 0.5 × TBE buffer and photographed over a UV light source (Kong et al. 2005).

### Linkage analysis

Data were analyzed using the chi-square ( $\chi^2$ ) test to ascertain goodness of fit between the expected ratio for a single dominant gene and the observed phenotypic segregation. Linkage analysis between the SSR or eSSR markers and the Hf-resistance genes, *H16* and *H17*, was performed with the software package MAPMAKER/EXP version 3.0 (Lander et al. 1987). Map units were computed by applying the Kosambi function (Kosambi 1944). The LOD score of 3 and the maximum distance of 50 cM were used in the determination of linkages.

## Results

### Hessian fly response phenotyping

BC<sub>1</sub>F<sub>1</sub> populations of Len/P921682 and Len/P921680 segregated, respectively, 55 resistant : 58 susceptible and 54 resistant : 49 susceptible. These numbers fit a 1:1 ratio ( $\chi^2 = 0.08$ ,  $P > 0.70$  and  $\chi^2 = 0.24$ ,  $P > 0.50$ , respectively) of a single dominant gene for Hessian fly resistance in both P921682 (*H16*) and P921680 (*H17*). The tests to phenotype the two BC<sub>1</sub>F<sub>1</sub> populations were definitive, all seedlings of the resistant parents P921682 (*H16H16*) and P921680 (*H17H17*), the durum donor lines PI 94587 and PI 428435, as well as F<sub>1</sub> seedlings derived from Len/P921682 and Len/P921680 were clearly not stunted and all plants of the susceptible parent line Len were clearly stunted.

### Molecular mapping of *H16* and *H17*


Because *H16* and *H17* were previously assigned to wheat chromosome 5A by segregation analysis, 40 5A-specific SSR markers were screened for linkage validation. However, no markers on either the long arm or short arm of chromosome 5A showed linkage relationships with Hf-resistance genes *H16* and *H17* (data not shown). Previously, the linkage

relationships with *H9* and/or *H10* suggested that *H16* and *H17* occupied a single linkage block (Obanni et al. 1988; Patterson et al. 1988; Ohm et al. 1995), and recently *H9* and *H10* were both placed on the distal region of wheat chromosome 1AS by molecular mapping (Kong et al. 2005; Liu et al. 2005a). Therefore, *H16* and *H17* were assumed to occupy the linkage block on chromosome 1A along with other Hf-resistance genes such as *H9* and *H10*. To test the hypothesis that *H16* and *H17* are on chromosome 1A, 62 SSR markers from chromosome 1A and 21 eSSR markers from group 1 (Table 1) were screened for potential linkage to *H16* or *H17*. As expected, no SSR markers other than a few from the short arm of 1A, and no eSSR markers other than two from 1AS showed linkage with *H16* and *H17*, which confirmed that both *H16* and *H17* are on the short arm of chromosome 1A. BC<sub>1</sub>F<sub>1</sub> populations of both *H16* and *H17* were further genotyped with additional SSR and eSSR markers based on BSA. Seven out of 62 SSR and 21 eSSR markers, including Cfa2153, Psp2999, Wem6b, Barc263, Wem3a, Wmc329 and Wmc24, showed linkage with *H16* and *H17*. Recombination analysis indicated that Hf-resistance gene *H16* was flanked by *Xpsp2999* and *Xwem6b/Xbarc263* with map distances of 3.7 and 5.5 cM, respectively. No recombination was observed between *Xbarc263* (Fig. 1) and *Xwem6b* in the Len\*2/P921682 population. The *Xcfa2153* marker locus was located distal to *H16* with 9.3 cM on chromosome 1AS. *Xwem3a*, *Xwmc329* and *Xwmc24* are proximal to *H16* at 11.1, 12.0, and 19.6 cM, respectively (Fig. 2a).

The *H16*-linked SSR and eSSR markers were also linked with *H17* on chromosome 1AS (Fig. 2b). *H17* is linked to the flanking markers *Xwem6b* and *Xpsp2999* with genetic distances of 5.1 and 6.2 cM, respectively. Marker *Xcfa2153* is distal to *H17* at 10.2 cM. Two other markers, *Xbarc263* and *Xwmc24*, are proximal to *H17* at 7.1, 17.3 cM, respectively. The eSSR marker *Xwem3a* co-segregated with *Xwmc329* and both are proximal to *H17* at 11.1 cM (Fig. 2b).

All of these seven linked markers were subsequently used to screen the resistant parent lines, P921682 and P921680, and the corresponding original durum donor lines PI 94587 and PI 428435 to determine the durum chromosomal segments transferred from the donors. A small terminal





**Fig. 1** DNA bands amplified from parents and 22 BC<sub>1</sub>F<sub>1</sub> plants derived from P921682(*H16H16*)/Len (*h16 h16*) with microsatellite primer pair *Xbarc263*. Lane 1 = 20-bp DNA ladder, Lane 2 = Len, Lane 3 = P921682, Lanes 4–25 = BC<sub>1</sub>F<sub>1</sub> plants (heterozygous resistant and homozygous susceptible are indicated by R and S, respectively). BC<sub>1</sub>F<sub>1</sub> plants in lanes 5 and


17 are recombinant between *Xbarc263* and *H16*. DNA fragments slightly larger than 210 and 240 bp were amplified, respectively, from the resistant parent and resistant BC<sub>1</sub>F<sub>1</sub> plants, versus the susceptible parent and susceptible BC<sub>1</sub>F<sub>1</sub> plants. These bands are indicated by the *black arrow* and *white arrow* on the left, respectively

chromosomal segment in wheat line P921680 carrying *H17* was found to be most likely transferred from the donor parent PI 428435 on the basis that *H17*-linked SSR markers *Xpsp2999* and *Xcfa2153*, distal to *H17*, and *Xwem6b* and *Xbarc263*, proximal to *H17* (the black region of chromosome 1AS in Fig. 2b) were not polymorphic between the wheat line P921680 (*H17H17*) and the durum *H17* donor, PI 428435. However, polymorphisms were detected at other proximal linked markers, *Xwem3a*, *Xwmc329*, and *Xwmc24* (the white region of chromosome 1AS in Fig. 2b). Interestingly, all of the seven *H16*-linked

markers showed polymorphisms between the wheat line P921682 (*H16H16*) and the durum *H16* donor, PI 94587. It is likely that a very small intercalary segment (the black region of chromosome 1AS in Fig. 2a) containing *H16* was transferred from PI 94587.


#### Physical mapping of the linked markers

DNA samples of CS and its derived deletion lines including del1AS-1 (FL 0.47) and del1AS3 (FL 0.86) were amplified using primer pairs of all of the *H16*- and



**Fig. 2** Linkage map of the short arm of wheat chromosome 1A showing the genetic map location of Hf-resistance gene *H16* (a), *H17* (b), and the consensus genetic map of the chromosomal region containing the *H16/H17* region on 1AS (c). Approximate distances in centi-Morgans (cM) and molecular markers are indicated on the left and the right, respectively. The letter X in front of each SSR locus name indicates the basic symbol for a molecular marker with

unknown function. The fraction lengths of the deletions are indicated in the parentheses, and the breakpoints of deletion bins are indicated to the right by arrows. The black regions of chromosome 1AS in (a) and (b) represent the durum donor-derived segment containing *H16* or *H17* in wheat lines P921682 and P921680, respectively. White regions of the chromosome in (a) and (b) represent the common wheat genetic backgrounds



**Fig. 3** DNA fragments amplified with simple-sequence-repeat primers Cfa2153 (a), Wem6b (b), Barc263 (c), and Wmc329 (d). Lane 1 = 20-bp DNA ladder, Lane 2 = CS del1AS-3 (0.86), Lane 3 = CS del1AS-1 (0.47), Lane 4 = CS, Lane 5 = P921680

(H17H17), Lane 6 = P921682 (H16H16), Lane 7 = Len, Lane 8 = 20-bp DNA ladder. Cfa2153, Wem6b, Barc263, and Wmc329 produced the specific fragments of 195, 155, 235, and 125 bp, respectively (pointed by arrows), only with CS

H17-linked SSR and eSSR markers including *Xcfa2153*, *Xpsp2999*, *Xwem6b*, *Xbarc263*, *Xwem3a*, *Xwmc329*, and *Xwmc24* to determine the physical locations of these linked markers. Each primer pair amplified DNA fragments of the expected size (s) from CS, but no corresponding fragments were amplified from CS del1AS-1 (FL 0.47) and CS del1AS-3 (FL 0.86) (Fig. 3a–d). This observation indicated that all of these linked markers along with Hf-resistance genes (*H16* and *H17*) are located in the distal 14% region on 1AS beyond the breakpoint of CS del1AS-3 (FL 0.86) (Figs. 2 and 3).

## Discussion

### Chromosome locations of Hf-resistance genes *H16* and *H17*

Recent molecular mapping analyses have provided conclusive evidence that Hf-resistance genes *H9*, *H10*, and *H11* are placed in the distal region of chromosome 1AS (Kong et al. 2005; Liu et al. 2005a). In this study, *H16* and *H17* are both mapped to the wheat chromosome 1AS by molecular techniques. Further evidence of Hf-resistance genes including *H9*, *H10*, *H11*, *H16*, and *H17* being located on chromosome 1AS came from the results of PCR amplification obtained from two CS deletion lines del1AS-1 (FL 0.47) and del1AS-3 (FL 0.86) with the primers for the closely linked SSR and eSSR markers in the current study and the

previous studies (Kong et al. 2005; Liu et al. 2005a; Peng and Lapitan 2005). Hf-resistant wheat lines P921682 and P921680 carry the dominant Hf-resistance gene *H16* or *H17*, respectively, which were initially localized on chromosome 5A by segregation analysis, showing linkage to *H9* and/or *H10* (Obanni et al. 1988; Patterson et al. 1988). *H9* was previously assigned to chromosome 5A based on linkage to *H6* (Stebbins et al. 1980), and *H6* was in turn mapped to 5A by monosomic analysis (Gallun and Patterson 1977). The error in placing gene *H6* on chromosome 5A, rather than on 1A by monosomic analysis could be the result of any of a number of factors including misidentification of monosomic genetic stocks. However, mapping of *H6*, originally and erroneously on chromosome 5A, emphasizes the importance of correctly mapping the first gene when it is used subsequently as a reference point for mapping additional genes. The original location of *H6* was not validated and all subsequent locations to the same region were based on close linkage to *H6*.

Linkage analysis and physical mapping of the linked molecular markers near the *H16* and *H17* loci positioned both of these Hf-resistance genes distal to the breakpoint of del3AS-3 (FL 0.86), or the distal 14% of the short arm of wheat chromosome 1A (Fig. 2a and b). Comparative analysis of *H16*/*H17* linkage maps and the consensus genetic map of 1AS (Somers et al. 2004; Paillard et al. 2003; Song et al. 2005; Bougot et al. 2002) is illustrated in Fig. 2c. In general, the marker order near *H16*/*H17* is similar to



that of the consensus genetic map on 1AS; the linkage region was flanked by *Xgwm136* and *Xwmc24* with genetic distance about 36 cM. Since the *H16/H17* loci are placed in the distal bin of 1AS (del1AS-3), we attempted to generate additional eSSR markers within the telomeric region of 1AS to define the physical end of the linkage map using the EST database. Of 21 eSSRs developed from the short arm of group 1 chromosomes (1AS, 1BS, and 1DS), only two eSSR markers, *Xwem6b* and *Xwem3a*, mapped proximal to *H16/H17* on 1AS (Fig. 2). The genetic locations of *Xwem6b* and *Xwem3a* reported in this study are in agreement with the locations revealed by eSSR physical mapping (Peng and Lapitan 2005). Out of 15 eSSR markers derived from either 1B or 1D, none shows linkage with *H16* or *H17*, which also supports the observation that *H16* and *H17* reside on 1AS.

DNA markers have facilitated the precise mapping of alleles or homoeo-loci. In this study, the map positions of *H16* and *H17* on chromosome 1AS are very similar (Fig. 2). However, the most closely linked flanking markers, *Xpsp2999* and *Xwem6b*, showed different banding patterns between Hf-resistant parent lines P921682 (*H16*) and P921680 (*H17*). Moreover, all of the more closely linked markers showed polymorphisms between the durum donors PI 94587 (*H16*) and PI 428435 (*H17*) except for the markers *Xwem3a*, *Xwmc24*, and *Xcfa2153* flanking this linkage region of 1AS (Fig. 2). It is noteworthy that, in our previous study, SSR markers *Xgwm136* and *Xgdm33* both are tightly linked distally to Hf-resistance gene *H9* at 1.7 and 2.2 cM, respectively (Kong et al. 2005). A similar mapping result with *H9* was reported by Liu et al. (2005a). However, in this study, neither *Xgwm136* nor *Xgdm33* showed linkage with *H16* or *H17*. It is unknown if *H9* is distal or proximal to *H16/H17* on 1AS. The actual linkage relationship between *H16/H17* and *Xgwm136/Xgdm33* cannot be observed due to the limited polymorphisms between Hf-resistant parent lines P921682 and P921680 and susceptible parent line Len in this gene-rich region on 1AS.

Comparison of Fig. 2a and b might raise the question whether *H16* and *H17* are at the same locus. However, analysis of the testcross: D6647 (susceptible to Hf)/IN80164 (*H16H16*)/PI 428435 (*H17H17*), all three parent lines being durum

wheat, showed that the genetic recombination frequency between *H16* and *H17* was 25 cM (Ohm et al. 1995).

#### A resistance gene cluster in the gene-rich distal region of 1AS

There are various reports that resistance genes to different pests and pathogens are linked and located in clusters observed in wheat (McIntosh et al. 1995, 2003; Adhikari et al. 2004); rice (Sardesai et al. 2002), maize (Hulbert et al. 2001), tomato (Dickinson et al. 1993), and soybean (Ashfield et al. 1998; Bachman et al. 2001). The genomic region that contains *H9*, *H10*, *H11*, *H16*, and *H17* is also particularly rich in genes for resistance against fungal pathogens. For example, *Pm3* for resistance to wheat powdery mildew (incited by *Blumeria graminis*) was also mapped with RFLP marker *BCD1434* (Hartl et al. 1993; Ma et al. 1994) and co-segregated with SSR marker *Xpsp2999* (Bougot et al. 2002) in the distal region of the short arm of chromosome 1A. At least 10 alleles (*Pm3a* to *Pm3j*) were identified at this locus (Zeller and Hsam 1998). Our previous study also confirmed that the STS marker derived from powdery mildew resistance gene *Pm3* was linked to Hf-resistance gene *H9* at a genetic distance of 4.5 cM (Kong et al. 2005). Leaf rust resistance gene *Lr10*, effective against *Puccinia triticina* Eriks, was also mapped in the same chromosomal region on 1AS (Schachermayr et al. 1997; Guyot et al. 2004). In order to determine the resistance gene order in this gene-rich region on 1AS, we designed primers for both *Pm3* and *Lr10* based on the available sequences in the database. Unfortunately, the *STS-Pm3* and *STS-Lr10* markers did not show polymorphism between the parent lines in the current study. However, because most of the SSR markers on 1AS showed linkage with *Pm3*, *Lr10*, and Hf-resistance genes including *H9*, *H10*, *H11*, *H16*, and *H17* (Kong et al. 2005; Liu et al. 2005a; Somers et al. 2004; Bougot et al. 2002; Guyot et al. 2004; Schachermayr et al. 1997), all of these resistance genes appear to compose a resistance gene cluster in the distal gene-rich region of 1AS.

The presence of multiple disease and insect resistance genes in this chromosomal region and positional cloning for both *Lr10* and *Pm3b* in bread wheat (Stein et al. 2000; Feullet et al. 2003; Yahiaoui

et al. 2004) make this genomic region much more attractive for future study including possible map-based cloning of any of the Hf-resistance genes in this region.

Marker-assisted selection being advanced by molecular technologies

Conventional wheat breeding for Hf-resistance relies on phenotypic selection through bioassays. More efficient and rapid breeding and deployment of Hf-resistant wheat varieties are critically needed because of the rapid evolution of Hf virulence. It has been postulated that the durability of resistance can be increased up to 20-fold by developing cultivars that contain multiple Hf-resistance genes (Gould 1986). Efficient pyramiding of effective Hessian fly resistance genes is possible with the aid of markers that co-segregate with the resistance genes, since expression of one gene effectively masks the presence of additional genes (Williams et al. 2003). The limiting factor for gene stacking is the lack of molecular markers specific to individual Hf-resistance genes. The results of this study are of practical significance to Hf resistance breeding. The specific and diagnostic SSR/eSSR markers closely linked to *H16* and *H17* identified in this study not only can assist wheat breeders in making parental selection but also will facilitate combining the Hf-resistance genes into elite breeding lines during cultivar development. These markers plus the other already mapped markers (Ma et al. 1994; Dweikat et al. 1997, 2002; Seo et al. 1997; Williams et al. 2003; Kong et al. 2005; Liu et al. 2005a, b, c; Sardesai et al. 2005; Wang et al. 2006; Zhao et al. 2006) will speed the development of breeding lines containing multiple resistance genes to develop broad-spectrum and durable resistance. For example, in this study, the two flanking markers, *Xpsp2999* and *Xwem6b*, are linked to *H16* at 3.7 and 5.5 cM, respectively (Fig. 2). The recombination frequencies (RF) between Hf-resistance gene *H16* and SSR markers *Xpsp2999*, *Xwem6b* are 3.6% and 5.2% (Haldane mapping function, Haldane 1919), respectively. The two RFs for *Xpsp2999* and *Xwem6b* translate into selection accuracies of 96.6% and 94.8%, respectively, if both of the markers are used separately. However, according to the product rule of the probability, the

selection accuracy will increase to nearly 100% [ $1 - (3.6\% \times 5.2\%)$ ] when these two flanking markers are used together.

**Acknowledgements** The authors acknowledge the financial support from USDA-CSREES Coordinated Agricultural Project (CAP) grant 2006-55606-16629, USDA-ARS and Purdue University. We also are grateful to Drs B. S. Gill, Kansas State University for providing seeds of wheat deletion lines of 1AS, and Junhua Peng, Colorado State University for identifying EST-derived microsatellites in the wheat EST database.

## References

- Anderson KG, Harris MO (2006) Does *R* gene resistance allow wheat to prevent plant growth effects associated with Hessian fly (Diptera: Cecidomyiidae) attack? J Econom Entomol 99:1842–1853
- Adhikari TB, Wallwork H, Goodwin SB (2004) Microsatellite markers linked to the *Stb2* and *Stb3* genes for resistance to *Septoria tritici* blotch in wheat. Crop Sci 44:1403–1411
- Ashfield T, Danzer JR, Held D, Clayton K, Keim P, Saghai Maroof MA, Webb DM, Innes RW (1998) *Rpg1*, a soybean gene effective against races of bacterial blight, maps to a cluster of previously identified disease resistance genes. Theor Appl Genet 96:1013–1021
- Berzonsky WA, Ding H, Haley SD, Harris MO, Lamb RJ, McKenzie RIH, Ohm HW, Patterson FL, Peairs F, Porter DR, Ratcliffe RH, Shanower TG (2003) Breeding wheat for resistance to insects. Plant Breed Rev 22:221–296
- Bachman MS, Tamulonis JP, Nickell CD, Bent AF (2001) Molecular markers linked to brown stem rot resistance genes, *Rbs1* and *Rbs2*, in soybean. Crop Sci 41:527–535
- Bougot Y, Lemonine J, Pavoine MT, Barloy D, Doussinault G (2002) Identification of a microsatellite marker associated with *Pm3* resistance alleles to powdery mildew in wheat. Plant Breed 121:325–329
- Cartwright WB, Caldwell RM, Compton LE (1959) Response of resistant and susceptible wheats to Hessian fly attack. Agron J 51:529–531
- Cox TS, Hatchett JH (1986) Genetic model for wheat Hessian fly (Diptera: Cecidomyiidae) interactions: strategies for deployment of resistance genes in wheat cultivars. Environ Entomol 15:24–31
- Delaney DE, Friebe BR, Hatchett JH, Gill BS, Hulbert SH (1995) Target mapping of rye chromatin in wheat by representational difference analysis. Genome 38:458–466
- Delibes A, Delmoral FJ, Martinsanchez JA, Meijias A, Gallego M, Casado D, Sin E, Lopezbrana I (1997) Hessian fly resistance gene transferred from chromosome 4M (V) of *Aegilops ventricosa* to *Triticum aestivum*. Theor Appl Genet 94:858–864
- Devos KM, Bryan GJ, Gale MD (1995) Application of two microsatellite sequences in wheat storage proteins as molecular markers. Theor Appl Genet 90:247–252
- Dickinson MJ, Jones DA, Jones JDG (1993) Close linkage between *CF2/Cf5* and *mi* resistance loci in tomato. Mol Plant-Microbe Interact 6:341–347

- Dweikat I, Ohm H, Patterson F, Cambron S (1997) Identification of RAPD markers for 11 Hessian fly resistance genes in wheat. *Theor Appl Genet* 94:419–423
- Dweikat I, Zhang W, Ohm H (2002) Development of STS markers linked to Hessian fly resistance gene *H6* in wheat. *Theor Appl Genet* 105:766–770
- Feuillet C, Treavella S, Stein N, Albar L, Nublat A, Keller B (2003) Map-based isolation of the leaf rust disease resistance gene *Lr10* from the hexaploid wheat (*Triticum aestivum* L.) genome. *Proc Natl Acad Sci USA* 100:15253–15258
- Gagné RJ, Hatchett JH (1989) Instars of the Hessian fly (Diptera: Cecidomyiidae). *Ann Entomol Soc Am* 82:73–79
- Gallun RL (1977) Genetic basis of Hessian fly epidemics. *Ann NY Acad Sci* 287:222–229
- Gallun RL, Deay HO, Cartwright WB (1961) Four races of Hessian fly selected and developed from an Indiana population. *Purdue Univ Agric Exp Stn Res Bull* 732:1–8
- Gallun RL, Hatchett JH (1968) Interrelationship between races of Hessian fly, *Mayetiola destructor* Say, and resistance in wheat. In *Proc Int Wheat Genet Symp*, 3rd, Canberra, Australia. 5–9 August. *Aust Acad Sci, Canberra, Australia*, pp 258–262
- Gallun RL, Patterson FL (1977) Monosomic analysis of wheat for resistance to Hessian fly. *J Hered* 68:223–226
- Gill BS, Hatchett JH, Raupp WJ (1987) Chromosomal mapping of Hessian fly-resistance gene *H13* in the D genome of wheat. *J Hered* 78:97–100
- Giovanini MP, Puthoff DP, Nemacheck JA, Mittapalli O, Saltzman KD, Ohm HW, Shukle RH, Williams CE (2006) Gene-to-gene defense of wheat against the Hessian fly lacks a classical oxidative burst. *Mol Plant Microbe Interact* 19:1023–1033
- Giovanini MP, Saltzman KD, Puthoff DP, Gonzalo M, Ohm HW, Williams CE (2007). A novel wheat gene encoding a putative chitin-binding lectin is associated with resistance against Hessian fly. *Molecular Plant Path* 8:69–82
- Gould F (1986) Simulation models for predicting durability of insect-resistant germ plasm: Hessian fly (Diptera: Cecidomyiidae)-resistant winter wheat. *Environ Entomol* 15:11–23
- Grover PBJ (1995) Hypersensitive response of wheat to Hessian fly (Diptera: Cecidomyiidae) biotypes on resistant wheat. *Environ Entomol* 18:687–690
- Gupta PK, Balyan HS, Edwards KJ, Isaac P, Korzun V, Röder MS, Gautier MF, Joudrier P, Schlatter AR, Dubcovsky J, De la Pena RC, Khairallah M, Penner G, Hayden MJ, Sharp P, Keller B, Wang RCC, Hardouin JP, Jack P, Leroy P (2002) Genetic mapping of 66 new microsatellite (SSR) loci in bread wheat. *Theor Appl Genet* 105:413–422
- Guyot R, Yahiaoui N, Feuillet C, Keller B (2004) In silico comparative analysis reveals a mosaic conservation of genes within a novel colinear region in wheat chromosome 1AS and rice chromosome 5S. *Funct Integr Genom* 4:47–58
- Haldane JBS (1919) The combination of linkage values, and the calculation of distances between the loci of linked factors. *J Genet* 8:299–309
- Harris MO, Freeman TP, Rohfritsch O, Anderson KG, Payne SA, Moore JA (2006) Virulent Hessian fly (Diptera: Cecidomyiidae) larvae induce a nutritive tissue during compatible interactions with wheat. *Ann Entomol Soc Am* 99:305–316
- Harris MO, Stuart JJ, Mohan M, Nair S, Lamb RJ, Rohfritsch O (2003) Grasses and gall midges: plant defense and insect adaptation. *Annu Rev Entomol* 48:549–577
- Hartl L, Weiss H, Zeller FJ, Jahoor A (1993) Use of RFLP markers for the identification of alleles of the *Pm3* locus conferring powdery mildew resistance in wheat (*Triticum aestivum* L.). *Theor Appl Genet* 86:959–963
- Hatchett JH, Gallun RL (1970) Genetics of the ability of the Hessian fly, *Mayetiola destructor*, to survive on wheats having different genes for resistance. *Ann Entomol Soc Am* 63:1400–1407
- Hulbert SH, Webb CA, Smith SM, Sun Q (2001) Resistance gene complexes: evolution and utilization. *Annu Rev Phytopathol* 39:285–312
- Kong L, Ohm HW, Cambron SE, Williams CE (2005) Molecular mapping determines that Hessian fly resistance gene *H9* is located on chromosome 1AS of wheat. *Plant Breed* 124:525–531
- Kosambi DD (1944) The estimation of map distances from recombination values. *Ann Eug* 12:172–175
- Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newburg L (1987) Mapmaker: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1:174–181
- Liu XM, Fritz AK, Reese JC, Wilde GE, Gill BS, Chen MS (2005a) *H9*, *H10*, and *H11* compose a cluster of Hessian fly resistance genes in the distal gene-rich region of wheat chromosome 1AS. *Theor Appl Genet* 110:143–148
- Liu XM, Brown-Guedira GL, Hatchett JH, Owuochi JO, Chen MS (2005b) Genetic characterization and molecular mapping of a Hessian fly resistance gene (*Hdic*) transferred from *T. turgidum* ssp. *Dicoccum* to common wheat. *Theor Appl Genet* 111:1308–1315
- Liu XM, Gill BS, Chen MS (2005c) Hessian fly resistance gene *H13* is mapped to a distal cluster of resistance genes in chromosome 6DS of wheat. *Theor Appl Genet* 111:243–249
- Ma ZQ, Sorrells ME, Tanksley SD (1994) RFLP markers linked to powdery mildew resistance genes *Pm1*, *Pm2*, *Pm3*, and *Pm4* in wheat. *Genome* 37:871–875
- Maas FB, Patterson FL, Foster JE, Ohm HW (1989) Expression and inheritance of resistance of ELS6404-160 durum wheat to Hessian fly. *Crop Sci* 29:23–28
- Martín-Sánchez JA, Gómez-Colmenarejo M, Del Moral J, Sin E, Montes MJ, González-Belinchón C, López-Braña I, Delibes A (2003) A new Hessian fly resistance gene (*H30*) transferred from the wild grass *Aegilops triuncialis* to hexaploid wheat. *Theor Appl Genet* 106:1248–1255
- McIntosh RA, Hart GE, Devos KM, Morris CF, Rogers WJ (2003) Catalogue of gene symbols for wheat, V. In: Raupp WJ (ed) *Annual wheat newsletter*, vol. 49. Kansas Agric Expt Sta, Manhattan, KS, USA
- McIntosh RA, Wellings CR, Park RF (1995) *Wheat rusts: an atlas of resistance genes*. CSIRO Publications, PO Box 89, East Melbourne, Australia
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in

- specific genomic regions by using segregating populations. *Proc Natl Acad Sci USA* 88:9828–9832
- Obanni M, Patterson FL, Foster JE, Ohm HW (1988) Genetic analysis of resistance of durum wheat PI 428435 to Hessian fly. *Crop Sci* 28:223–226
- Ohm HW, Sharma HC, Patterson FL, Ratcliffe RH, Obanni M (1995) Linkage relationships among genes on wheat chromosome 5A that condition resistance to Hessian fly. *Crop Sci* 35:1603–1607
- Paillard S, Schnurbusch T, Winzeler M, Messmer M, Sourdille P, Abderhalden O, Keller B, Schachermayr G (2003) An integrative genetic linkage map of winter wheat (*Triticum aestivum* L.). *Theor Appl Genet* 107:1235–1242
- Patterson FL, Foster JE, Ohm HW (1988) Gene *H16* in wheat for resistance to Hessian fly. *Crop Sci* 28:652–654
- Patterson FL, Gallun RL (1977) Linkage in wheat of the *H3* and *H9* genetic factors for resistance to Hessian fly. *J Hered* 68:293–296
- Peng JH, Lapitan NLV (2005) Characterization of EST-derived microsatellites in the wheat genome and development of eSSR markers. *Funct Integr Genom* 5:80–96
- Pestsova E, Ganai MW, Röder MS (2000) Isolation and mapping of microsatellite markers specific for the D genome of bread wheat. *Genome* 43:689–697
- Ratcliffe RH, Hatchett JH (1997) Biology and genetics of the Hessian fly and resistance in wheat. In: Bondari K (ed) *New developments in entomology: Research signpost*, Trivandrum, India, pp 47–56
- Robinson RJ, Miller BS, Miller HL, Mussman HC, Johnson JA, Jones ET (1960) Chloroplast number in leaves of normal wheat plants and those infested with Hessian fly or treated with maleic hydrazine. *J Econom Entomol* 53:560–566
- Röder MS, Korzun V, Wendehake K, Plaschke J, Tixier MH, Leroy P, Ganai MW (1998) A microsatellite map of wheat. *Genetics* 149:2007–2023
- Saghai-Maroo MA, Soliman KM, Jorgensen RA, Allard RW (1984) Ribosomal DNA spacer length polymorphism in barley: Mendelian inheritance chromosomal location and population dynamics. *Proc Natl Acad Sci USA* 81:8014–8018
- Sardesai N, Kumar A, Rajyashri KR, Nair S (2002) Identification and mapping of an AFLP marker linked to *Gm7*, a gall midge resistance gene and its conversion to a SCAR marker for its utility in marker aided selection in rice. *Theor Appl Genet* 105:691–698
- Sardesai N, Nemacheck JA, Subramanyam S, Williams CE (2005) Identification and mapping of *H32*, a new wheat gene conferring resistance to Hessian fly. *Theor Appl Genet* 111:1167–1173
- Schachermayr G, Feuillet C, Keller B (1997) Molecular markers for the detection of the wheat leaf rust resistance gene *Lr10* in diverse genetic backgrounds. *Mol Breed* 3:65–74
- Seo YW, Johnson JW, Jarret RL (1997) A molecular marker associated with the *H21* Hessian fly resistance gene in wheat. *Mol Breed* 3:177–181
- Smith CM, Quisenberry SS, du Toit F (1999) The value of conserved wheat germplasm possessing arthropod resistance. In: Clement SL, Quisenberry SS (eds) *Global plant genetic resources for insect-resistant crops*. CRC Press, Boca Raton, pp 25–49
- Somers DJ, Isaac P, Edwards K (2004) A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* 109:1105–1114
- Song QJ, Shi JR, Singh S, Fickus EW, Costa JM, Lewis L, Gill BS, Ward R, Cregan BP (2005) Development and mapping of microsatellite (SSR) markers in wheat. *Theor Appl Genet* 110:550–560
- Stebbins NB, Patterson FL, Gallun RL (1982) Interrelationships among genes *H3*, *H6* and *H9* for Hessian fly resistance. *Crop Sci* 22:1029–1032
- Stebbins NB, Patterson FL, Gallun RL (1980) Interrelationships among wheat genes for resistance to Hessian fly. *Crop Sci* 20:177–180
- Stein N, Feuillet C, Wicker T, Schlagenhauf E, Keller B (2000) Subgenome chromosome walking in wheat: a 450-kb physical contig in *Triticum monococcum* L. spans the *Lr10* resistance locus in hexaploid wheat (*Triticum aestivum* L.). *Proc Natl Acad Sci USA* 97:13436–13441
- Stephenson P, Bryan G, Kirby J, Collins A, Devos K, Busso C, Gale M (1998) Fifty new microsatellite loci for the wheat genetic map. *Theor Appl Genet* 97:946–949
- Wang T, Xu SS, Harris MO, Hu J, Liu L, Cai X (2006) Genetic characterization and molecular mapping of Hessian fly resistance genes derived from *Aegilops tauschii* in synthetic wheat. *Theor Appl Genet* 113:611–618
- Williams CE, Collier CC, Nemacheck JA, Chengzhi L, Cambron SE (2002) A lectin-like wheat gene responds systematically to attempted feeding by avirulent first-instar Hessian fly larvae. *J Chem Ecol* 28:1411–1428
- Williams CE, Collier CC, Sardesai N, Ohm HW, Cambron SE (2003) Phenotypic assessment and mapped markers for *H31*, a new wheat gene conferring resistance to Hessian fly (Diptera: Cecidomyiidae). *Theor Appl Genet* 107:1516–1523
- Yahiaoui N, Srichumpa P, Dudler R, Keller B (2004) Genome analysis at different ploidy levels allows cloning of the powdery mildew resistance gene *Pm3b* from hexaploid wheat. *Plant J* 37:528–538
- Zeller FJ, Hsam SLK (1998) Progress in breeding for resistance to powdery mildew in common wheat (*Triticum aestivum* L.). In: Slinkard AE (ed) *Proc 9th Int Wheat Genet Symp*, Saskatoon, Saskatchewan, Canada, vol 1, pp 178–180
- Zhao HX, XM Liu, Chen MS (2006) *H22*, a major resistance gene to the Hessian fly (*Mayetiola destructor*), is mapped to the distal region of wheat chromosome 1DS. *Theor Appl Genet* 113:1491–1496